arXiv:math/0503002v1 [math.CO] 1 Mar 2005

Counting formulas associated with some random matrix averages

Peter J. Forrester* and Alex Gamburd†
*Department of Mathematics and Statistics, University of Melbourne,
Victoria 3010, Australia ;
† Department of Mathematics, University of California, Santa Cruz
and Department of Mathematics, Stanford University,
Stanford, CA 94305 USA

Abstract: Moments of secular and inverse secular coefficients, averaged over random matrices from classical groups, are related to the enumeration of non-negative matrices with prescribed row and column sums. Similar random matrix averages are related to certain configurations of vicious random walkers and to the enumeration of plane partitions. The combinatorial meaning of the average of the characteristic polynomial of random Hermitian and Wishart matrices is also investigated, and consequently several simple universality results are derived.

1 Introduction

The richness of random matrix theory was greatly enhanced during the last couple of years of the millennium by the discovery of its intimate connections with increasing subsequences and non-intersecting lattice paths in enumerative and asymptotic combinatorics (for reviews see [1, 13]). These topics relate to non-negative integer matrices via the celebrated Robinson-Schensted-Knuth (RSK) correspondence [15, 20]. A special class of non-negative matrices are so called magic squares — they have the property that the sum of the elements in any row or column is equal to a prescribed positive integer j say. Such matrices are natural objects in RSK theory, so one might expect a relationship between magic squares and random matrices. This is indeed the case, and its development forms the main theme of the recent work [8] (see [6] for related results). One of the goals of the present paper is to continue the development of this theme.

Magic squares are special cases of non-negative integer $m \times n$ rectangular matrices in which the sum of the elements in each row j is equal to μ_j, while the sum of the elements in each column k is equal to $\tilde{\mu}_k$. Without loss of generality, for the purpose of enumeration we can insist that $\mu_1 \geq \cdots \geq \mu_m$ and $\tilde{\mu}_1 \geq \cdots \geq \tilde{\mu}_n$ so that $\mu := (\mu_1, \ldots, \mu_m)$ and $\tilde{\mu} := (\tilde{\mu}_1, \ldots, \tilde{\mu}_n)$ form partitions. We denote the total number of such matrices by $N_{\mu\tilde{\mu}}$. Writing the partitions in terms of the frequencies of their parts by

$$\mu = \langle 1^{a_1} \cdots l^{a_l} \rangle, \qquad \tilde{\mu} = \langle 1^{b_1} \cdots l^{b_l} \rangle \tag{1.1}$$

where $l = \max(\mu_1, \tilde{\mu}_1)$, it was proved in [8, Thm. 2] that for $N \ge \max(\sum_{j=1}^l j a_j, \sum_{j=1}^l j b_j)$

$$N_{\mu\tilde{\mu}} = \mathbb{E}_{M \in U(N)} \prod_{j=1}^{l} (\mathrm{Sc}_j(M))^{a_j} \overline{(\mathrm{Sc}_j(M))}^{b_j}. \tag{1.2}$$

In (1.2) the average is over matrices M chosen from the group $U(N)$ at random with respect to the Haar measure (uniform distribution), while $\mathrm{Sc}_j(M)$ is the jth secular coefficient of the characteristic polynomial of M,

$$P_M(z) = \det(M - zI_N) = \sum_{j=0}^{N} \mathrm{Sc}_j(M)(-z)^{N-j}. \tag{1.3}$$

From (1.2) it follows that the number $H_k(j)$ of $k \times k$ magic squares, specified as $k \times k$ non-negative integer matrices with the sum of elements in each row and column equal to j, is for $N \ge kj$, given by the formula

$$H_k(j) = \mathbb{E}_{M \in U(N)} |\mathrm{Sc}_j(M)|^{2k}. \tag{1.4}$$

We will extend the formula (1.2) in a number of directions. The first relates to counting formulas analogous to (1.2) in the case that the matrices are constrained by a symmetry property. Two such formulas, both relating to $n \times n$ symmetric matrices, are known from [8]. Thus with the row sums (which must be equal to the column sums as the matrices are symmetric) labelled by the partition $\mu = (\mu_1, \dots, \mu_n) = \langle 1^{a_1} \cdots l^{a_l} \rangle$ we have that for $|\mu| := \sum_{j=1}^l j a_j$ even, $N \ge |\mu|$, and all elements on the diagonal zero, the total number N_μ^{O} of such non-negative integer matrices is given by

$$N_\mu^{\mathrm{O}} = \mathbb{E}_{M \in O(N)} \prod_{j=1}^{l} (\mathrm{Sc}_j(M))^{a_j}. \tag{1.5}$$

If instead all elements on the diagonal are permitted to be even, the total number N_μ^{Sp} of such matrices is given by

$$N_\mu^{\mathrm{Sp}} = \mathbb{E}_{M \in USp(2N)} \prod_{j=1}^{l} (\mathrm{Sc}_j(M))^{a_j}. \tag{1.6}$$

In (1.5) the average is over the group of orthogonal matrices $O(N)$, while in (1.6) it is over the group of $2N \times 2N$ unitary symplectic matrices $USp(2N)$ (both with the Haar measure). In Section 2 we will extend (1.5) and (1.6) to the case that the diagonal elements have no special restriction. We also give analogous formulas in the case of square matrices symmetric about both the diagonal and anti-diagonal, and for matrices symmetric about the centre point of the matrix. That such symmetrizations relate to averages over classical groups is a consequence of results of Rains [24], and Baik and Rains [2]. Another tractable case in terms of a random matrix average to be considered is the setting relating to (1.2) but with the entries of the matrix restricted to 0's and 1's. If the matrix has a 2×2 block structure, with the two diagonal blocks having non-negative integer entries, and the off diagonal blocks having entries 0 or 1, counting formulas of the type (1.2) can be obtained by studying ratios of characteristic polynomials averaged over the classical groups. This is done in Section 3.

The topic of Section 4 is the relationship between enumeration formulas (1.2) and (1.6) and certain classes of non-intersecting lattice paths, or equivalently certain configurations of vicious random walkers. This is motivated by the graphical representation of the RSK correspondence in terms of non-intersecting lattice paths [17, 14]. In Section 5 we describe connections between moments of characteristic polynomials and enumeration of certain classes of plane partitions. Finally, in Section 6, we revisit the question of the combinatorial meaning of the expected value of characteristic polynomials of random complex Hermitian Wigner matrices, and take up a similar study in relation to Wishart matrices.

2 Further symmetrizations of the square

The RSK correspondence gives a bijection between weighted $n \times n$ non-negative integer matrices, entries x_{ij} weighted by $(\alpha_i\beta_j)^{x_{ij}}$, and pairs of weighted semi-standard tableaux of content n. In the latter one member of the pair is weighted by

$$\alpha_1^{\#1's}\alpha_2^{\#2's}\cdots\alpha_n^{\#n's}, \tag{2.1}$$

while the other member is weighted by (2.1) but with the α's replaced by the β's (the exponents in (2.1) are determined by the entries of the matrix). Summing (2.1) over all allowed semi-standard tableaux of a given shape κ gives the combinatorial definition of the Schur polynomial, and this way one obtains for the generating function of the weighted matrices the identity

$$\frac{1}{\prod_{i,j=1}^{n}(1-\alpha_i\beta_j)} = \sum_{\kappa} s_\kappa(\alpha_1,\dots,\alpha_n)s_\kappa(\beta_1,\dots,\beta_n). \tag{2.2}$$

This is well known in the theory of the Schur polynomial, and is called the Cauchy formula (see e.g. [22]). We remark that the case of $m \times n$ matrices ($m < n$ say for definiteness) follows by simply setting $\alpha_{m+1} = \cdots = \alpha_n = 0$.

The fact that the entries x_{ij} are weighted by $(\alpha_i\beta_j)^{x_{ij}}$ tells us the coefficient of

$$\alpha_1^{\eta_1}\cdots\alpha_n^{\eta_n}\beta_1^{\rho_1}\cdots\beta_n^{\rho_n}$$

in (2.2) counts the number of matrices for which the sum of the elements in row i equals η_i, while the sum of the elements in row j equals ρ_j. Because (2.2) is symmetric in the α_i's and the β_j's, without loss of generality we can restrict attention to the case that $\eta_1 \ge \cdots \ge \eta_n$, $\rho_1 \ge \cdots \ge \rho_n$ and thus η and ρ form partitions, to be denoted μ and $\tilde{\mu}$ respectively say. The task is then to extract the coefficient of $\alpha^\mu\beta^{\tilde{\mu}}$ (in an obvious multivariable shorthand notation).

Only terms on the RHS of (2.2) with $|\kappa| = |\mu| = |\tilde{\mu}|$ can contribute to the coefficient of $\alpha^\mu\beta^{\tilde{\mu}}$. Because of this, we can use a result of Baik and Rains [2] expressing the RHS of (2.2) with the largest part of κ restricted to be no greater than N as an average over $U(N)$,

$$\sum_{\kappa:\kappa_1\le N} s_\kappa(\alpha_1,\dots,\alpha_n)s_\kappa(\beta_1,\dots,\beta_n) = \mathbb{E}_{U\in U(N)}\prod_{j=1}^{n}\det(I_N+\alpha_j\bar{U})\det(I_N+\beta_j U). \tag{2.3}$$

In particular with μ, $\tilde{\mu}$ specified as in (1.1), (2.3) agrees with (2.2) at the required order provided

$$N \geq |\mu| = \sum_{j=1}^{l} j a_j = \sum_{j=1}^{l} j b_j. \tag{2.4}$$

The extraction of the sought coefficient from (2.3) is immediate, and we reclaim (1.2), provided N obeys the inequality (2.4).

We remark in passing, that since we have [5, 8]

$$\mathbb{E}_{U \in U(N)} \prod_{j=1}^{m} \det(I_N + \alpha_j U) \prod_{j=1}^{n} \det(I_N + \overline{U}/\beta_j) = \frac{1}{(\beta_1 \ldots \beta_n)^N} s_{N^n}(\alpha_1, \ldots, \alpha_m; \beta_1, \ldots, \beta_n), \tag{2.5}$$

(in fact this equation will be of independent interest below) where N^n denotes the partition with n parts all equal to N, it is also true that $N_{\mu\tilde{\mu}}$ is equal to the coefficient of $\alpha^{\mu}\beta^{N-\tilde{\mu}}$ in the Schur function $S_{N^n}(\{\alpha_j, \beta_j\}_{j=1,\ldots,n})$, provided N is large enough as required by (2.4).

From the present perspective, to extend the enumeration formula (1.2) to classes of symmetrized non-negative matrices we require a formula analogous to (2.3) for the corresponding partial generating function. Indeed such a formula (due to Littlewood [21]) is known for symmetric matrices with all elements zero on the diagonal,

$$\sum_{\lambda' \,\mathrm{even}} s_\lambda(\alpha_1, \ldots, \alpha_n) = \prod_{1 \leq i < j \leq n} \frac{1}{1 - \alpha_i \alpha_j}, \tag{2.6}$$

and for symmetric matrices with all elements even on the diagonal,

$$\sum_{\lambda \,\mathrm{even}} s_\lambda(\alpha_1, \ldots, \alpha_n) = \prod_{1 \leq i \leq j \leq n} \frac{1}{1 - \alpha_i \alpha_j}. \tag{2.7}$$

Combining (2.6) with

$$\mathbb{E}_{M \in O(N)} \prod_{j=1}^{n} \det(I_N + \alpha_j M) = \sum_{\substack{l(\lambda) \leq N \\ \lambda' \,\mathrm{even}}} s_\lambda(\alpha_1, \ldots, \alpha_n) \tag{2.8}$$

and (2.7) with

$$\mathbb{E}_{M \in USp(2N)} \prod_{j=1}^{n} \det(I_{2N} + \alpha_j M) = \sum_{\substack{\lambda_1 \leq 2N \\ \lambda \,\mathrm{even}}} s_\lambda(\alpha_1, \ldots, \alpha_n) \tag{2.9}$$

allows us to reclaim (1.5) and (1.6) respectively. The finitizations (2.8) and (2.9) of Littlewood's formula are due to Rains [24].

Formulas analogous to (2.3) are also known in the case of symmetric matrices with no special constraint on the diagonal, matrices symmetric about both the diagonal and anti-diagonal, and matrices with a point reflection symmetry about the centre. Let us first consider symmetric matrices with no special constraint on the diagonal.

For a symmetric matrix, the row sum equals the column sum so we can take for the weights $\alpha_i \beta_j = \sqrt{q_i q_j}$ and extract the coefficient of q^{μ} to read off the number of non-negative integer matrices with row sums equal to μ. Since each off diagonal element in position ij of a symmetric matrix is identical to the element in position ji, we can equivalently restrict attention to the lower triangular portion $i \leq j$ of the matrix, and weight the elements in strictly lower triangular positions $i < j$ by $q_i q_j$, and the diagonal elements by q_i. The generating function for weighted symmetric matrices is thus seen to be

$$\frac{1}{\prod_{i=1}^{n}(1-q_i)\prod_{i<j}(1-q_iq_j)} \tag{2.10}$$

and we seek the coefficient of q^μ in this expression. According to an identity of Littlewood [21] we have that (2.10) is equal to

$$\sum_{\kappa} s_\kappa(q_1,\dots,q_n). \tag{2.11}$$

It has been shown by Baik and Rains [2] that (2.11) constrained by the size of the largest part of κ_1 can be expressed as a random matrix average,

$$\sum_{\kappa:\kappa_1\leq N} s_\kappa(q_1,\dots,q_n) = \mathbb{E}_{U\in O(N)} \det(I_N+U)\prod_{j=1}^{n}\det(I_N+q_jU). \tag{2.12}$$

With knowledge of the above results, we can now easily obtain an enumeration formula for symmetric non-negative integer matrices with prescribed row sums.

Proposition 1. *Consider the set of all symmetric $n\times n$ non-negative integer matrices. Label the row sums (or equivalently column sums) by the partition*

$$\mu = (\mu_1,\dots,\mu_n) = \langle 1^{a_1}\cdots l^{a_l}\rangle.$$

For $N \geq |\mu|$ the total number $N_\mu^{\boxslash}$ of such matrices is given as a random matrix average by

$$N_\mu^{\boxslash} = \mathbb{E}_{M\in O(N)}\det(I_N+M)\prod_{j=1}^{l}(\mathrm{Sc}_j(M))^{a_j} \tag{2.13}$$

(cf. (1.5)).

Proof. Only terms in (2.11) with $|\kappa| = |\mu|$ can contribute to the coefficient of q^μ in (2.10), so for $N \geq |\mu|$ the latter is the same as the coefficient of q^μ in (2.12). Extracting the coefficient gives (2.13). $\square$

We remark that since $\det(I_N+M)=0$ for $M\in O^-(N)$ (because $\lambda=-1$ is an eigenvalue) we can restrict the matrices M in (2.13) to $M\in O^+(N)$ (the matrices U in (2.12) can similarly be restricted).

Next, we note that the results of Baik and Rains in fact allow the diagonal sum to be prescribed. The relevant generating function is [2]

$$\frac{1}{\prod_{i=1}^{n}(1-\alpha q_i)\prod_{i<j}(1-q_iq_j)} = \sum_{\kappa}\alpha^{\sum_{j=1}^{n}(-1)^{j-1}\kappa_j}s_\kappa(q_1,\dots,q_n) \tag{2.14}$$

and for $\kappa_1 \leq N$ this sum can be written as the random matrix average

$$\mathbb{E}_{U\in O(N)} \det(I_N + \alpha U) \prod_{j=1}^{n} \det(I_N + q_j U). \tag{2.15}$$

Hence, with the diagonal sum prescribed to be equal to p say and the total number of symmetric non-negative matrices now denoted $N^{\boxslash}_{\mu,p}$, the enumeration formula (2.13) should be modified to read

$$N^{\boxslash}_{\mu,p} = \mathbb{E}_{M\in O(N)}\mathrm{Sc}_p(M)) \prod_{j=1}^{l} (\mathrm{Sc}_j(M))^{a_j}. \tag{2.16}$$

We now turn our attention to the task of enumerating according to row sums $2n \times 2n$ non-negative integer matrices $[x_{i,j}]_{i,j=1,\dots,2n}$ with a reflection symmetry about the diagonal, $x_{i,j} = x_{j,i}$ $(i > j)$, and about the anti-diagonal, $x_{i,j} = x_{i,2n+1-j}$ $(i > 2n + 1 - j)$. The generating function for weighted symmetric matrices of this type is

$$\prod_{i=1}^{n} \frac{(1 + \chi_1 q_i)}{(1 - \chi_0 q_i)} \frac{1}{\prod_{i,j=1}^{n}(1 - q_i q_j)} \tag{2.17}$$

where $\chi_0 = 1$ $(\chi_1 = 1)$ in the case that the elements on the diagonal (anti-diagonal) are unrestricted, while $\chi_0 = 0$ $(\chi_1 = 0)$ in the case the elements on the diagonal are restricted to be zero (anti-diagonal are restricted to be even). We know from [2] that the random matrix average

$$\mathbb{E}_{U\in U(N)} \frac{\det(1 + \chi_0 U)}{\det(I_N - \chi_1 U)} \prod_{j=1}^{n} \det(I_N + q_j U) \det(I_N + q_j \bar{U}) \tag{2.18}$$

correctly reproduces (2.17) up to and including terms $O(q_j^{2N+1})$ in the q_j. (In the case $\chi_1 = 1$, (2.18) is to be interpreted as the limiting value for $\chi_1 \to 1^-$.) The following enumeration result is now evident.

Proposition 2. *Consider the set of all symmetric $2n \times 2n$ non-negative integer matrices which have the further constraint of being symmetric about the anti-diagonal. Introduce possible constraints on the diagonal and anti-diagonal elements according to the values of χ_0 and χ_1 noted below (2.17). With the row sums labelled by μ, we have that for $N \geq \frac{1}{2}|\mu|$ the number of matrices in the set, $N^{\boxtimes}_{\mu}$ say, is given by the coefficient of q^{μ} in the random matrix average (2.18).*

As our next example of a symmetry constraint, we turn our attention to the case of $2n \times 2n$ matrices invariant with respect to reflections in the point $(n+1/2, n+1/2)$ (here we are thinking of the matrix as labelled by a grid of $2n \times 2n$ lattice points $\{(i,j) : 1 \leq i,j \leq 2n\}$). For such matrices $x_{i,j} = x_{2n+1-i,2n+1-j}$. The generating function for this class of matrices is

$$\frac{1}{\prod_{i,j=1}^{n}(1 - q_i q_j)^2}. \tag{2.19}$$

It follows from (2.2) and (2.3) that up and including terms $O(q_i^N)$ (2.19) is equal to

$$\Big(\mathbb{E}_{U\in U(N)} \prod_{j=1}^{n} \det(I_N + q_j \bar{U}) \det(I_N + q_j U)\Big)^2. \tag{2.20}$$

We thus have the following enumeration result.

Proposition 3. *Consider the set of all* $2n \times 2n$ *non-negative integer matrices with the point reflection symmetry* $x_{i,j} = x_{2n+1-i,2n+1-j}$. *With the row sums labelled by* μ, *we have that for* $N \ge |\mu|$, *the number of matrices in the set* $N_\mu^{\boxdot}$ *say is given by the coefficient of* q^μ *in the random matrix average (2.20).*

As another extension of (1.2) we consider not a symmetry constraint on the matrix, but rather a restriction on the entries of the matrix. These we take to be either 0 or 1. For $m \times n$ matrices of this type, weighted by $(\alpha_i\beta_j)^{x_{ij}}$, the generating function is

$$\prod_{i=1}^{m}\prod_{j=1}^{n}(1+\alpha_i\beta_j). \tag{2.21}$$

According to the dual Cauchy identity,

$$\prod_{i,j=1}^{n}(1+\alpha_i\beta_j) = \sum_{\mu} s_{\mu'}(\alpha_1,\dots,\alpha_n)s_\mu(\beta_1,\dots,\beta_n), \tag{2.22}$$

where μ' denotes the partition conjugate to μ (see e.g. [22]). Analogous to (2.3), we have that this sum restricted by the size of the largest part of μ can be written as a random matrix average

$$\sum_{\mu:\mu_1\le N} s_{\mu'}(\alpha_1,\dots,\alpha_n)s_\mu(\beta_1,\dots,\beta_n) = \mathbb{E}_{U\in U(N)}\prod_{j=1}^{n}\frac{\det(I_N+\alpha_jU)}{\det(I_N-\beta_j\bar{U})}. \tag{2.23}$$

Because of the occurrence of the dual partition in (2.22), μ_1 cannot exceed n for a non-zero contribution and so (2.23) is independent of N for $N \ge n$.

We want to extract from this the coefficient of $\alpha^\mu\beta^{\tilde{\mu}}$. For this purpose we introduce inverse secular coefficients $\mathrm{Rc}_p(U)$ according to

$$\frac{1}{\det(I_N-xU)} = \sum_{p=0}^{\infty}x^p\mathrm{Rc}_p(U).$$

In terms of the eigenvalues $e^{i\theta_j}$ $(j=1,\dots,N)$ of U we have that

$$\mathrm{Rc}_p(U) = h_p(e^{i\theta_1},\dots,e^{i\theta_N})$$

where h_p denotes the pth complete symmetric function (see [22]). Making use too of (1.3) we obtain for the sought enumeration the following result.

Proposition 4. *Consider the set of all* $m\times n$ *matrices in which the entries take on the values 0 or 1. Prescribe the row sums and column sums by the partitions* μ *and* $\tilde{\mu}$ *respectively, which are to be written in terms of the frequency of their parts by (1.1). For* $N \ge \min(m,n,\sum_{j=1}^{l} ja_j, \sum_{j=1}^{l} jb_j)$ *the total number* $N_{\mu\tilde{\mu}}^{0,1}$ *of such matrices is given in terms of a random matrix average by*

$$N_{\mu\tilde{\mu}}^{0,1} = \mathbb{E}_{U\in U(N)}\prod_{j=1}^{l}(\mathrm{Sc}_j(U))^{a_j}(\mathrm{Rc}_j(\bar{U}))^{b_j}. \tag{2.24}$$

We remark that for (2.24) to be non-zero we require $\mu_1 \le m$, $\tilde{\mu}_1 \le n$ and $|\mu| = |\tilde{\mu}|$.

3 Block matrix structures and ratios of characteristic polynomials

We have seen in the previous section how generating functions for non-negative integer matrices, and 0-1 matrices, are related to Schur function identities which in turn are related to averages over the classical groups. Results obtained in [4, 2] tell us this strategy can also be carried through for classes of block matrices which relate to averages over the unitary, symplectic and orthogonal groups.

In relation to the unitary group, let the block structured $(k_1+l_1)\times(k_2+l_2)$ non-negative integer matrix

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} \tag{3.1}$$

be such that A is a $k_1\times k_2$ non-negative integer matrix; D is a $l_1\times l_2$ non-negative integer matrix; B is a $k_1\times l_2$ 0-1 matrix; and C is a $l_1\times k_2$ 0-1 matrix. Suppose the entries of the block $X=[x_{ij}]$ are weighted by $g_{ij}^{x_{ij}}$ with

$$g_{ij} = \begin{cases} \alpha_i\gamma_j & \text{for} \quad X=A \\ \beta_i\delta_j & \text{for} \quad X=D \\ \alpha_i\delta_j & \text{for} \quad X=B \\ \beta_i\gamma_j & \text{for} \quad X=C. \end{cases} \tag{3.2}$$

Such matrices have for their generating function

$$\prod_{i=1}^{k_1}\prod_{j=1}^{k_2}\frac{1}{1-\alpha_i\gamma_j}\prod_{i=1}^{l_1}\prod_{j=1}^{l_2}\frac{1}{1-\beta_i\delta_j}\prod_{i=1}^{k_1}\prod_{j=1}^{l_2}(1+\alpha_i\delta_j)\prod_{i=1}^{l_1}\prod_{j=1}^{k_2}(1+\beta_i\gamma_j),$$

thus containing as special cases both the LHS of (2.2) and (2.21). The coefficient of $\alpha^\mu\beta^{\tilde{\mu}}\gamma^\nu\delta^{\tilde{\nu}}$ tells us the number of matrices (3.1) with prescribed row and column sums. Moreover generalizations of both the Cauchy identities (2.2), (2.22), and their finitizations (2.3), (2.23) are known [4, 2], allowing for the counting function to be expressed as a random matrix average.

To state these generalizations requires introducing the functions [22]

$$\mathrm{HS}_\lambda(\alpha_1,\dots,\alpha_k;\beta_1,\dots\beta_l) = \det(a_{\lambda_i+j-1})_{1\le i,j\le l(\lambda)}, \tag{3.3}$$

where a_k denotes the coefficient of x^k in

$$\frac{\prod_{j=1}^{l}(1+\beta_j x)}{\prod_{i=1}^{k}(1-\alpha_i x)}.$$

The HS_λ are referred to as the hook Schur functions. In terms of this the generalization of the Cauchy identities is

$$\sum_\lambda \mathrm{HS}_\lambda(\alpha_1,\dots,\alpha_{k_1};\beta_1,\dots\beta_{l_1})\mathrm{HS}_\lambda(\gamma_1,\dots,\gamma_{k_2};\delta_1,\dots\delta_{l_2})$$
$$=\prod_{i=1}^{k_1}\prod_{j=1}^{k_2}\frac{1}{1-\alpha_i\gamma_j}\prod_{i=1}^{l_1}\prod_{j=1}^{l_2}\frac{1}{1-\beta_i\delta_j}\prod_{i=1}^{k_1}\prod_{j=1}^{l_2}(1+\alpha_i\delta_j)\prod_{i=1}^{l_1}\prod_{j=1}^{k_2}(1+\beta_i\gamma_j) \tag{3.4}$$

while the generalizations of their finitizations is

$$\mathbb{E}_{U\in U(N)} \frac{\prod_{i=1}^{k_1}\det(I_N+\alpha_i U)\prod_{j=1}^{k_2}\det(I_N+\gamma_j\bar{U})}{\prod_{m=1}^{l_1}\det(I_N-\beta_m U)\prod_{n=1}^{l_2}\det(I_N-\delta_n\bar{U})} = \sum_{\lambda_1\le N} \mathrm{HS}_\lambda(\alpha_1,\dots,\alpha_{k_1};\beta_1,\dots\beta_{l_1})\mathrm{HS}_\lambda(\gamma_1,\dots,\gamma_{k_2};\delta_1,\dots\delta_{l_2}). \tag{3.5}$$

Arguing as in the derivation of the counting formula (1.2) given in the first three paragraphs of Section 2, the following generalization of (1.2) and (2.24) is immediate.

Proposition 5. *Let* $\mathbf{a}=(a_1,\dots,a_{k_1})$, $\mathbf{b}=(b_1,\dots,b_{l_1})$, $\mathbf{c}=(c_1,\dots,c_{k_2})$, $\mathbf{d}=(d_1,\dots,d_{l_2})$, *where the* a_j, b_j, c_j, d_j *are nonnegative integers. From these arrays form partitions*

$$\mu=\langle 1^{a_1}\cdots k_1^{a_{k_1}}\rangle,\ \ \tilde{\mu}=\langle 1^{b_1}\cdots l_1^{b_{l_1}}\rangle,\ \ \nu=\langle 1^{c_1}\cdots k_2^{c_{k_2}}\rangle,\ \ \tilde{\nu}=\langle 1^{d_1}\cdots l_2^{d_{l_2}}\rangle.$$

Let $N_{\mu\tilde{\mu}\nu\tilde{\nu}}$ *denote the number of matrices (3.1) with*

$$\mathrm{row}(A,B)=\mu,\ \ \mathrm{row}(C,D)=\nu,\ \ \mathrm{col}(A,C)=\tilde{\mu},\ \ \mathrm{col}(B,D)=\tilde{\nu}$$

where the notation row(X,Y) *refers to the row sums across* X *and* Y *given they are horizontal neighbors in a block matrix, and col*(X,Y) *refers to the column sums down* X *and* Y *given they vertical neighbors in a block matrix.*

For

$$N\ge\max\left(\sum_1^{k_1} ja_j,\sum_1^{l_1} jb_j,\sum_1^{k_2} jac_j,\sum_1^{l_2} jd_j,\right)$$

we have

$$\mathbb{E}_{U_N}\prod_{i=1}^{k_1}(\mathrm{Sc}_i(M))^{a_i}\prod_{j=1}^{l_1}\overline{(\mathrm{Sc}_j(M))}^{b_j}\prod_{m=1}^{k_2}(\mathrm{Rc}_m(M))^{c_m}\prod_{n=1}^{l_2}\overline{(\mathrm{Rc}_n(M))}^{d_n}=N_{\mu\tilde{\mu}\nu\tilde{\nu}}. \tag{3.6}$$

As reviewed in the Introduction, a relationship with averages over the orthogonal and symplectic groups comes about when the matrix is constrained to be symmetric. Thus in (3.1) we must take

$$k_1=k_2=k,\quad l_1=l_2=l,\quad A=A^T,\quad D=D^T,\quad C=B^T. \tag{3.7}$$

Correspondingly, for $i\ne j$, the weights (3.2) are to be replaced by

$$g_{ij}=\begin{cases}\sqrt{\alpha_i\alpha_j} & \text{for}\quad X=A\\ \sqrt{\beta_i\beta_j} & \text{for}\quad X=D\\ \sqrt{\alpha_i\beta_j} & \text{for}\quad X=B\\ \sqrt{\beta_i\alpha_j} & \text{for}\quad X=C.\end{cases} \tag{3.8}$$

As in the meaning of (1.5) and (1.6), the constraint on the diagonal elements will determine whether the relationship is with an average over the symplectic group, or an average over the orthogonal group. It turns out that the constraint relevant to the symplectic (orthogonal) group is that all elements on the diagonal of A be even (zero), while those on D be zero (even). The corresponding generating functions are therefore given by

$$\begin{aligned} G^{\mathrm{Sp}}(\{\alpha_i\},\{\beta_j\}) &= \prod_{1\le i\le j\le k}\frac{1}{1-\alpha_i\alpha_j}\prod_{1\le i<j\le l}\frac{1}{1-\beta_i\beta_j}\prod_{i=1}^{k}\prod_{j=i}^{l}(1+\alpha_i\beta_j) \\ G^{\mathrm{O}}(\{\alpha_i\},\{\beta_j\}) &= \prod_{1\le i<j\le k}\frac{1}{1-\alpha_i\alpha_j}\prod_{1\le i\le j\le l}\frac{1}{1-\beta_i\beta_j}\prod_{i=1}^{k}\prod_{j=i}^{l}(1+\alpha_i\beta_j) \end{aligned}$$

In relation to these generating functions, one has as generalizations of the Littlewood identities (2.6) and (2.7) [4]

$$\begin{aligned} \sum_{\lambda\ \mathrm{even}} \mathrm{HS}_\lambda(\alpha_1,\dots\alpha_k;\beta_1,\dots,\beta_l) &= G^{\mathrm{Sp}}(\{\alpha_i\},\{\beta_j\}) \\ \sum_{\lambda'\ \mathrm{even}} \mathrm{HS}_\lambda(\alpha_1,\dots\alpha_k;\beta_1,\dots,\beta_l) &= G^{\mathrm{O}}(\{\alpha_i\},\{\beta_j\}). \end{aligned}$$

And with the largest part of the partitions in the sum restricted, the LHS's have the finitizations [2]

$$\mathbb{E}_{U\in USp(2N)}\frac{\prod_{i=1}^{k}\det(I_{2N}+\alpha_iU)}{\prod_{j=1}^{l}\det(I_{2N}-\beta_jU)} = \sum_{\substack{\lambda_1\le 2N\\ \lambda\ \mathrm{even}}} \mathrm{HS}_\lambda(\alpha_1,\dots\alpha_k;\beta_1,\dots,\beta_l) \tag{3.9}$$

$$\mathbb{E}_{U\in O(N)}\frac{\prod_{i=1}^{k}\det(I_{N}+\alpha_iU)}{\prod_{j=1}^{l}\det(I_{N}-\beta_jU)} = \sum_{\substack{l(\lambda)\le N\\ \lambda'\ \mathrm{even}}} \mathrm{HS}_\lambda(\alpha_1,\dots\alpha_k;\beta_1,\dots,\beta_l). \tag{3.10}$$

As a consequence, the following enumeration results hold.

Proposition 6. *Let* $\mathbf{a}=(a_1,\dots,a_k)$, $\mathbf{b}=(b_1,\dots,b_l)$, *where* a_j, b_j *are non-negative integers, and from these arrays form partitions*

$$\mu=\langle 1^{a_1}\cdots k^{a_k}\rangle, \qquad \nu=\langle 1^{b_1}\cdots l^{b_l}\rangle$$

Consider block matrices (3.1) with constraints (3.7) and the further constraint that all diagonal entries of A *are even (zero) while all those of* D *are zero (even). Let* $N_{\mu\nu}^{\mathrm{Sp}}$ $(N_{\mu\nu}^{\mathrm{O}})$ *denote the number of such matrices with*

$$\mathrm{row}(A,B)=\mathrm{col}(A,C)=\mu, \qquad \mathrm{row}(C,D)=\mathrm{col}(B,D)=\nu.$$

We have

$$\begin{aligned} N_{\mu\nu}^{\mathrm{Sp}} &= \mathbb{E}_{M\in USp(2N)}\prod_{i=1}^{k}(\mathrm{Sc}_i(M))^{a_i}\prod_{j=1}^{l}(\mathrm{Rc}_j(M))^{c_j} \\ N_{\mu\nu}^{\mathrm{O}} &= \mathbb{E}_{M\in O(N)}\prod_{i=1}^{k}(\mathrm{Sc}_i(M))^{a_i}\prod_{j=1}^{l}(\mathrm{Rc}_j(M))^{c_j}. \end{aligned}$$



Figure 1: Example of non-intersecting lattice paths corresponding to returning lock step vicious walkers. For the first (last) n steps the walkers must move to the right (left) or stay stationary. The weight of the configuration shown here is $\alpha_1^2\alpha_2^2\alpha_3^2\alpha_4\beta_1^3\beta_2\beta_3\beta_4^2$.

4 Relationship to non-intersecting paths

The purpose of this section is to relate the random unitary matrix average in (2.3) and the unitary symplectic average in (2.9) to configurations of weighted non-intersecting lattice paths. This then allows us to give combinatorial interpretations to (1.2) and (1.6) relating to non-intersecting lattice paths rather than integer matrices. That such interpretation are possible can be anticipated from the RSK correspondence: the semi-standard tableaux therein have a well known interpretation in terms of weighted non-intersecting lattice paths (see e.g. [25]). However neither in the case of (2.3) nor (2.9) will our non-intersecting paths correspond to the conventional ones related to semi-standard tableaux, and so a separate discussion is warranted.

In relation to (2.3), mark points on the x-axis at $x = 1, \dots, N$. Move each point to the line $y = 1$ according to the rule that each x coordinate must either stay the same (weight unity) or increase by one (weight α_1), with the proviso that all x coordinates must remain distinct. Connect the points between $y = 0$ and $y = 1$ by segments, which must either be vertical (weight unity), or right diagonal (weight α_1). Repeat this procedure a total of n times, with each right diagonal segment at step j weighted by α_j.

After step n, perform another n steps, but now with the segments either vertical (weight unity) or left diagonal (weight β_{2n+1-j} in step $n + j$). The segments again must not intersect, and are further conditioned to return along $y = 2n$ to the same x coordinates ($x = 1, \dots, N$) as they began (see Figure 1 for an example). The resulting non-intersecting lattice paths are equivalent to a special case of the lock-step model of vicious random walkers [9, 10]. For general initial positions along $y = 0$ ($l_1^{(0)}, \dots, l_N^{(0)}$ say), and final positions along $y = 2n$ ($l_1, \dots, l_N$ say), the generating function G_{2n} for the weighted paths can be written as an $N \times N$ determinant according to

$$G_{2n}(l_1^{(0)}, \dots, l_N^{(0)}; l_1, \dots, l_N) = \det \Big[g_{2n}(l_j^{(0)}; l_k) \Big]_{j,k=1,\dots,N} \tag{4.1}$$

where

$$g_{2n}(l^{(0)}; l) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \prod_{j=1}^{n} (1 + \alpha_j e^{-i\theta_j})(1 + \beta_j e^{i\theta_j}) e^{-i(l - l^{(0)})\theta} \, d\theta. \tag{4.2}$$

From the well known identity

$$\det \Big[\frac{1}{2\pi} \int_{-\pi}^{\pi} h(\theta) e^{-i(j-k)\theta} \, d\theta \Big]_{j,k=1,\dots,N} = \Big\langle \prod_{j=1}^{N} h(\theta_j) \Big\rangle_{U(N)} \tag{4.3}$$

we see that in the case of interest (initial positions = final positions, all one unit apart), we have

$$G_{2n}(l_1^{(0)}, \dots, l_N^{(0)}; l_1, \dots, l_N) \Big|_{\substack{l_j^{(0)} = l_j = j \\ (j=1,\dots,N)}} = \mathbb{E}_{U \in U(N)} \prod_{j=1}^{n} \det(I_N + \alpha_j \bar{U}) \det(I_N + \beta_j U). \tag{4.4}$$

The following interpretation of the average over $U(N)$ in (1.2) is now evident.

Proposition 7. *Consider the set of all non-intersecting paths of the type depicted in Figure 1. Impose the further constraint that the number of right diagonal segments between $y = j - 1$ and $y = j$ ($j = 1, \dots, n$) is equal to μ_j, and the total number of left diagonal segments between $y = 2n - j$ and $y = 2n - j + 1$ ($j = 1, \dots, n$) is equal to $\tilde{\mu}_j$. The total number of such lattice paths is given by the RHS of (1.2).*

We note that in the setting of Proposition 7 at most $|\mu| = \sum_{j=1}^{n} \mu_j$ (which must equal $|\tilde{\mu}|$) different walkers, counted to the left from $x = N$, can move. Thus we see immediately that (1.2) must be independent of N for $N \geq |\mu|$.

Let us now turn our attention to the non-intersecting lattice path interpretation of (2.9). For this mark points on the x-axis at $x = 1, \dots, N$. Also count steps $j = 1, \dots, 2n$ as the points are moved to the lines $y = 1, \dots, 2n$ in order, and the corresponding segments are drawn to connect the points. Let this process proceed by the rule that the lattice paths must not intersect, and that at odd numbered steps the segments must be either vertical or right diagonal, while at even numbered steps the segments must either be vertical or left diagonal. Weight the diagonal segments in steps $2j - 1, 2j$ by a_j, b_j respectively. It is further required that all points remain to the right of the y-axis, which can be thought of as a wall (see Figure 2 for an example). For general initial positions along $y = 0$ ($l_1^{(0)}, \dots, l_N^{(0)}$) and finishing positions along $y = 2n$ ($l_1, \dots, l_N$), all to the right of the wall, the generating function G_{2n}^{wall} for the weighted paths has



Figure 2: Example of non-intersecting lattice paths with initial (final) spacings along $y = 0$ ($y = 2n$) one unit apart starting at $x = 1$. All paths are restricted to $x \geq 1$, and segments can be vertical or right diagonal for odd steps, and vertical or left diagonal for even steps. The weight of the configuration shown here is $\alpha_1^3 \alpha_2^4 \alpha_3^2 \alpha_4^3$.

a determinant form [10]. In the special case that $\{a_i\} = \{b_i\} = \{\alpha_i\}$ (in any order), and that the initial and final positions are spaced one unit apart starting at $x = 1$, this determinant can be expressed as the random matrix average (2.9) [11], thus providing the following interpretation of the latter in terms of non-intersecting paths.

Proposition 8. *Consider the set of all non-intersecting lattice paths of the type depicted in Figure 2. Impose the further constraint that the sum of the number of right diagonal segments at step* $2j-1$ *and the number of left diagonal segments at step* $2j$ *is equal to* μ_j*. The total number of such paths is given by the RHS of (1.6).*

5 Relationship to plane partitions

In this section the random unitary matrix average (2.5), the random unitary symplectic average (2.9), and the random orthogonal average (2.15) will be related to the enumeration of certain classes of plane partitions [22, 26]. A plane partition $\mathcal{P}$ is a finite set of lattice points $\{(i, j, k)\} \subseteq \mathbb{N}^3$ with the property that if $(a, b, c) \in \mathcal{P}$ and $1 \leq i \leq a$, $1 \leq j \leq b$, $1 \leq k \leq c$, then $(i, j, k) \in \mathcal{P}$. A plane partition is symmetric if $(i, j, k) \in \mathcal{P}$ if and only if $(j, i, k) \in \mathcal{P}$. The height of stack

(i,j) is the largest value of k for which there exists a point (i,j,k) in the plane partition.

The study of plane partitions was initiated by MacMahon [23] who proved that the generating function for plane partitions fitting in the box

$$\mathcal{B}(a,b,c)=\{(i,j,k)|1\le i\le a,1\le j\le b,1\le k\le c\} \tag{5.1}$$

is given by

$$\prod_{i=1}^{a}\prod_{j=1}^{b}\prod_{k=1}^{c}\frac{1-q^{i+j+k-1}}{1-q^{i+j+k-2}}. \tag{5.2}$$

Taking the limit $q\to 1$ gives for the total number of plane partitions fitting inside $\mathcal{B}(a,b,c)$, $\#\mathcal{P}(a,b,c)$ say, the evaluation

$$\#\mathcal{P}(a,b,c)=\prod_{i=1}^{a}\prod_{j=1}^{b}\prod_{k=1}^{c}\frac{i+j+k-1}{i+j+k-2}. \tag{5.3}$$

We can express the generating function (5.2) for plane partitions in terms of Schur functions. Thus from the combinatorial definition of Schur functions $s_\lambda(x_1,\dots,x_n)$ as a sum over weighted semi-standard tableaux of shape λ and content $\{1,\dots,n\}$, it follows that

$$s_{b^a}(q^{a+c},q^{a+c-1},\dots,q)$$

where b^a denotes a partition with a parts all of which are equal to b, is the generating function for plane partitions strictly decreasing down columns with exactly a rows each of length b and with the largest stack of height less than or equal to $a+c$. The strictly decreasing constraint can be eased by removing $a-i+1$ from the boxes in row i to bijectively obtain a plane partition which is a subset of $\mathcal{B}(a,b,c)$. Consequently an alternative expression for (5.2) is given by

$$q^{-ba(a+1)/2}s_{b^a}(q^{a+c},q^{a+c-1},\dots,q). \tag{5.4}$$

Making use of (2.5) shows that for $a=n$, $c=m$ and $b=N$, (5.4) can be expressed as

$$\mathbb{E}_{U\in U(N)}\prod_{j=1}^{m}\det(I_N+\alpha_jU)\prod_{j=1}^{n}\det(I_N+\overline{U}/\beta_j)$$

with $\alpha_i=q^{n+i}$ and $\beta_i=q^i$. Taking the limit $q\to 1$ gives the sought relationship between (5.3) and a random matrix average.

Proposition 9. *Denote by $P_M^*(z)$ the (reciprocal) characteristic polynomial of $M\in U(N)$, so that $P_M(z)=\det(I_N-zM)$. For $|z|=1$ we have*

$$\#\mathcal{P}(a,b,c)=\mathbb{E}_{U\in U(c)}\Big(P_M^*(z)\Big)^b\Big(\overline{P_M^*(z)}\Big)^a. \tag{5.5}$$

We remark that in [27], the case $a=b$ of the RHS of (5.5) has been given a combinatorial interpretation involving two-rowed lexicographic arrays.

The random matrix average (5.5) is an example of a class of multi-dimensional integrals possessing gamma function evaluations (see e.g. [12]). This implies

$$\begin{aligned} \#\mathcal{P}(a,b,c) &= \prod_{j=0}^{c-1} \frac{\Gamma(a+b+1+j)\Gamma(2+j)}{\Gamma(a+1+j)\Gamma(b+1+j)} \\ &= \frac{G(1+a+b+c)}{G(1+a+b)} \frac{G(1+a)}{G(1+a+c)} \frac{G(1+b)}{G(1+b+c)} G(c+2) \end{aligned} \tag{5.6}$$

where $G(z)$ is the Barnes G-function, related to the gamma function by the functional equation $G(z+1)=\Gamma(z)G(z)$, $G(1)=1$. In contrast to the formula (5.3), the latter formula is well suited for asymptotic analysis.

Consider now symmetric plane partitions fitting inside the box (5.1) with $b=a$. The generating function for such plane partitions, with the additional constraint that the heights of all stacks on the diagonal are even and bounded by $2c$, has the product form [26]

$$\prod_{1\le i\le j\le a} \frac{1-q^{i+j+2c}}{1-q^{i+j}}. \tag{5.7}$$

Here only points on and above the diagonal are weighted. Denoting their total number by $\#\mathcal{P}_{\rm e}^{\rm sym}(a,2c)$, we see by taking the limit $q\to 1$ that [7]

$$\#\mathcal{P}_{\rm e}^{\rm sym}(a,2c) = \prod_{1\le i\le j\le a} \frac{i+j+2c}{i+j}. \tag{5.8}$$

The generating function (5.7) and thus counting formula (5.8) can be expressed in terms of Schur functions by making use of a bijection between tableaux and symmetrical plane partitions. Consider then semi-standard tableaux of content $\{1,\dots,a\}$. Suppose furthermore that each row is of even length, and the first row is constrained to be less than or equal to $2c$. From this construct the diagonal and upper triangular portion of a symmetrical plane partition by associating with grid points (i,j), $i\le j$, stacks of height

$$h_{i,j} = \#i's + \#(i+1)'s + \cdots + \#(a+i-j)'s \ \text{ in row } i \text{ of the tableaux.}$$

By weighting each square labelled j in the tableaux by q^j, we see that each stack at grid point (i,j) is weighted $q^{h_{i,j}}$, and furthermore on the diagonal this weight is q^{λ_i} where λ_i is the length of row i of the tableau. It follows that (5.7) can be expressed in terms of Schur functions according to

$$\sum_{\substack{\lambda\subseteq(2c)^a \\ \lambda\,\text{even}}} s_\lambda(q^a, q^{a-1},\dots,q). \tag{5.9}$$

Recalling (2.9) gives the sought relationship with an average over the unitary symplectic group.

Proposition 10. *Denote by* $P_M^{\rm Sp}(z)$ *the characteristic polynomial of* $M\in USp(2c)$ *so that* $P_M^{\rm Sp}(z)=\det(\lambda I_{2c}-M)$. *We have*

$$\#\mathcal{P}_{\rm e}^{\rm sym}(a,2c) = \mathbb{E}_{M\in USp(2c)}\Big(P_M^{\rm Sp}(-1)\Big)^a.$$

Analogous to (5.6), the above average is of a type which admits a gamma function evaluation (see e.g. [12]). This allows (5.8) to be written

$$\begin{aligned}
\#\mathcal{P}^{\rm sym}_{\rm e}(a,2c) &= 2^{2ca}\prod_{j=1}^{c}\frac{\Gamma(1+c+j)\Gamma(\frac{1}{2}+a+j)}{\Gamma(1+c+a+j)\Gamma(\frac{1}{2}+j)} \\
&= 2^{2ca}\frac{G(2+2c)}{G(2+c)}\frac{G(2+c+a)}{G(2+2c+a)}\frac{G(\frac{3}{2}+a+c)}{G(\frac{3}{2}+a)}\frac{G(\frac{3}{2})}{G(\frac{3}{2}+c)}.
\end{aligned}$$

Finally, consider the same weighted symmetric plane partitions as described in the paragraph below (5.8), but with stacks now restricted to heights $\le c$ and without the constraint that the stacks on the diagonal be even. The generating function is

$$\sum_{\lambda\subseteq c^a} s_\lambda(q^a,q^{a-1},\dots,q).$$

In an obvious notation, after taking the limit $q\to 1$ and recalling (2.12) we obtain

$$\#\mathcal{P}^{\rm sym}(a,c) = \mathbb{E}_{M\in O(c)}\Big(P_M^{\rm O*}(-1)\Big)^{a+1}.$$

6 Counting formulas associated with the characteristic polynomials of random Hermitian and Wishart matrices

Let X be an $n\times n$ Hermitian matrix. Let the diagonal elements x_{ii} be chosen independently according to a probability distribution $\mathcal{D}_1$, with the property that

$$\mathbb{E}_{\mathcal{D}_1} x_{ii} = 0. \tag{6.1}$$

Let the upper triangular elements x_{ij}, $i<j$, (which may be complex) be chosen independently according to a probability distribution $\mathcal{D}_2$, with the properties that

$$\mathbb{E}_{\mathcal{D}_2} x_{ij} = 0, \qquad \mathbb{E}_{\mathcal{D}_2}|x_{ij}|^2 = \sigma_2^2. \tag{6.2}$$

The following result, generalizing a number of results presented in [8][1], gives the expected value of the characteristic polynomial of X in terms of the classical Hermite polynomial $H_n(x)$.

Proposition 11. *Let X be Hermitian and specified in terms of $\mathcal{D}_1$ and $\mathcal{D}_2$ as above. With the monic rescaled Hermite polynomial specified in terms of the classical Hermite polynomial by*

$$h_n(x) = 2^{-n/2}H_n\Big(\frac{x}{\sqrt{2}}\Big)$$

we have

$$\mathbb{E}_{\mathcal{D}_1,\mathcal{D}_2}\det(\lambda I_N - X) = \sigma_2^N h_N\Big(\frac{\lambda}{\sigma_2}\Big). \tag{6.3}$$

---

[1] We take this opportunity to correct the statement of Theorem 15 in [8]. It should read: $\mathbb{E}_{\mu_N}(P_M^{2k}(x)) = h_N^{(k)}(x)$, where $h_N^{(k)}$ are orthogonal polynomials with respect to the weight $(t-x)^{2k}e^{-t^2}$.

Proof. By definition

$$\det(\lambda I_N - X) = \sum_{P \in S_N} \varepsilon(P) \prod_{l=1}^{N} (\lambda_{l,P(l)} - x_{l,P(l)}), \tag{6.4}$$

where $\varepsilon(P)$ denotes the parity of P and

$$\lambda_{i,j} = \begin{cases} \lambda & \text{if } i = j; \\ 0, & \text{if } i \neq j. \end{cases}$$

The specifications (6.1), (6.2) tell us that the only non-zero terms in (6.4) after averaging over $\mathcal{D}_1, \mathcal{D}_2$ are those for which P consists entirely of fixed points ($P(j) = j$) and 2-cycles ($P(j_1) = j_2$ and $P(j_2) = j_1$, $j_1 \neq j_2$). Let there then be $N - 2j$ fixed points and j 2-cycles. Such permutations have parity $(-1)^j$. Each fixed point contributes a factor λ, while each 2-cycle contributes σ_2^2. As the number of ways of choosing the $(N - 2j)$ fixed points and the j 2-cycles is

$$\binom{N}{2j} \frac{(2j)!}{2^j j!}$$

we see that

$$\mathbb{E}_{\mathcal{D}_1,\mathcal{D}_2} \det(\lambda I_N - X) = \sum_{j=0}^{[N/2]} (-1)^j \binom{N}{2j} \frac{(2j)!}{2^j j!} \lambda^{N-2j} \sigma_2^{2j}.$$

But this is precisely the power series expansion of the polynomial in question. □

Next we turn our attention to the mean characteristic polynomial of so called chiral matrices, that is matrices of the form

$$K := \begin{bmatrix} 0_{n \times n} & X_{n \times p} \\ (X^\dagger)_{p \times n} & 0_{p \times p} \end{bmatrix}, \tag{6.5}$$

where we require that $n \geq p$. These matrices have exactly $n - p$ zero eigenvalues, with the remaining $2p$ eigenvalues given by ± the positive square roots of the eigenvalues of the non-negative matrix $X^\dagger X$. We specify that the elements of X be chosen independently according to a probability distribution $\mathcal{D}$ with the properties

$$\mathbb{E}_{\mathcal{D}} x_{ij} = 0, \qquad \mathbb{E}_{\mathcal{D}} |x_{ij}|^2 = \sigma^2. \tag{6.6}$$

The analogue of Proposition 11 can readily be deduced.

Proposition 12. *Let K be a random chiral matrix as specified above. We have*

$$\mathbb{E}_{\mathcal{D}} \det(\lambda I_{n+p} - K) = p! \sigma^{2p} \lambda^{n-p} L_p^{n-p}((\lambda/\sigma)^2), \tag{6.7}$$

where $L_m^a(x)$ denotes the classical Laguerre polynomial.

Proof. Because of the first specification in (6.6), we see that in the analogue of (6.4) for $\det(\lambda I_{n+p} - K)$, the only terms after averaging will again result entirely from fixed points and 2-cycles. The fact that K has $n - p$ zero eigenvalues implies we require there be a minimum of

$n-p$ fixed points. Thus we must consider the cases that the number of fixed points is equal to $n+p-2j$ and the number of 2-cycles is equal to j for each $j=0,\dots,p$.

Not all permutations with these specifications give a non-zero contribution. For the latter, because of the zero blocks in (6.5), in relation to the 2-cycles $(j_1 j_2)$ we require

$$j_1 \in \{1,\dots,n\}, \qquad j_2 \in \{n+1,\dots,n+p\}.$$

Hence we must choose $(n-p)+(p-j)$ fixed points from the first of these sets and $(p-j)$ from the second. The number of distinct ways to do this is

$$\binom{n}{n-j}\binom{p}{p-j}.$$

The j 2-cycles can then be chosen in $j!$ different ways. Since again each fixed point contributes a factor λ, while each 2-cycle contributes a factor σ^2, we see that

$$\begin{aligned}\mathbb{E}_{\mathcal{D}} \det(\lambda I_{n+p} - K) &= \sigma^{2p}\lambda^{n-p}\sum_{j=0}^{p}\binom{n}{n-j}\binom{p}{p-j}j!\Big(\frac{\lambda}{\sigma}\Big)^{2(p-j)} \\ &= p!\sigma^{2p}\lambda^{n-p}\sum_{j=0}^{p}\binom{n}{p-j}\frac{1}{j!}\Big(\frac{\lambda}{\sigma}\Big)^{2j}.\end{aligned}$$

The sum in this expression is precisely $L_p^{n-p}((\lambda/\sigma)^2)$. □

Due to the relationship between the chiral matrices $X^\dagger X$ as noted below (6.5), we have the following result for the expected value of the characteristic polynomial for Wishart matrices.

Corollary 1. *Let X be a $n \times p$ matrix with elements independently chosen according to the distribution $\mathcal{D}$ with the properties (6.6). We have*

$$\mathbb{E}_{\mathcal{D}} \det(\lambda I_p - X^\dagger X) = p!\sigma^p L_p^{n-p}(\lambda/\sigma). \tag{6.8}$$

Acknowledgements

We thank the organizers of the program ‘Random matrix methods in number theory’, held at the Newton Institute during the first half of 2004, for inviting us to participate and so facilitating our collaboration. The work of PJF was supported by the Australian Research Council. The work of AG was supported in part by NSF Postdoctoral Fellowship.

References

[1] D. Aldous, P. Diaconis, *Longest increasing subsequences: from patience sorting to the Baik-Deift-Johansson theorem*, BAMS, **36**, 1999, 413-432.

[2] J. Baik and E. M. Rains, *Algebraic aspects of increasing subsequences,* Duke Mathematical Journal, **109**, 2001, 1-65.

[3] T. H. Baker and P. J. Forrester, *Finite-N fluctuation formulas for random matrices*, J. Stat. Phys., **88**, 1997, 1371-1386.

[4] A. Berele and J. B. Remmel, *Hook flag characters and their combinatorics,* J. Pure Appl. Algebra, **35**, 1985, 225-245.

[5] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein, and N. C. Snaith, *Autocorrelation of random matrix polynomials*, Communications in Mathematical Physics, **237**, 2003, 365-395.

[6] B. Conrey and A. Gamburd, *Pseudomoments of the Riemann zeta-function and pseudomagic squares*, to appear in Journal of Number Theory.

[7] M. De Sainte-Catherine and G. Viennot, *Enumeration of certain Young tableaux with bounded heights*, Lecture Notes in Mathematics, **1234**, 1986, 58-67.

[8] P. Diaconis and A. Gamburd, *Random matrices, magic squares and matching polynomials,* Electronic Journal of Combinatorics, **11**(2), 2004, #R2.

[9] M. E. Fisher, *Walks, walls, welting and melting,* J. Stat. Phys., **34**, 1984, 667-729.

[10] P. J. Forrester, *Probability of survival of vicious walkers near a cliff,* J. Phys. A., **22**, 1989, L609-L613.

[11] P. J. Forrester, *Random walks and random permutations,* J. Phys. A., **34**, 2001, L417-L423.

[12] P. J. Forrester, *Log-gases and Random Matrices*, www.ms.unimelb.edu.au/m̃atpjf/matpjf.html

[13] P. J. Forrester, *Growth models, random matrices and Painlevé transcendents,* Nonlinearity, **16**, 2003, R27–R49

[14] P. J. Forrester and E. M. Rains, *Interpretations of some parameter dependent generalizations of classical matrix ensembles,* Probab. Theory Relat. Fields, **131**, 2005, 1–61

[15] W. Fulton, *Young Tableaux,* London Mathematical Society, Student Texts **35**, Cambridge Univ. Press, 1997.

[16] J. Gravner, C. A. Tracy, H. Widom, *Limit theorems for height fluctuations in a class of discrete space and time growth models,* J. Stat. Phys., **102**, 2001, 1085-1132.

[17] K. Johansson, *Non-intersecting paths, random tilings and random matrices,* Probab. Theory Relat. Fields, **123**, 2002, 225-280.

[18] J. P. Keating and N. C. Snaith, *Random matrix theory and* $\zeta(\frac{1}{2}+it)$, Commun. Math. Phys., **214**, 2000, 57-89.

[19] J. P. Keating and N. C. Snaith, *Random matrix theory and L-functions at* $s=\frac{1}{2}$, Commun. Math. Phys., **214**, 2000, 91-110.

[20] D. E.E. Knuth, *Permutations, matrices, and generalized Young tableaux*, Pacific J. Math., **34**, 1970, 709-727.

[21] D. E. Littlewood, *The theory of group characters and matrix representations of groups,* Oxford at the Clarendon Press, 1940.

[22] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Second edition, Oxford University Press, New York, 1995.

[23] P. A. MacMahon, *Combinatory Analysis,* Cambridge University Press, 1915.

[24] E. M. Rains, *Increasing subsequences and the classical groups,* Electronic Journal of Combinatorics, **5**, 1998, #R12.

[25] B. E. Sagan, *The Symmetric Group*, 2nd edition, Springer, 2000.

[26] R. P. Stanley. *Enumerative Combinatorics, Vol. 2.* Cambridge University Press, 1999.

[27] E. Strahov. *Moments of characteristic polynomials enumerate two-rowed lexicographic arrays*, Electronic Journal of Combinatorics, **10**, 2003, #R24.